

Pfizer Oncology together™

INJECTION
Retacrit™
epoetin alfa-epbx
Pfizer Oncology

RETACRIT™ Billing and Coding Guide



Please see Important Safety Information and Indications on pages 14-16, and full Prescribing Information, including **BOXED WARNING** and Medication Guide, available at [RetacritHCP.com](https://www.RetacritHCP.com).

RETACRIT is a trademark of Pfizer Inc.

Pfizer Oncology Together is a trademark of Pfizer Inc.

Introduction

Pfizer Inc. has developed this reference guide to assist healthcare providers (HCPs) with understanding coding for RETACRIT (epoetin alfa-epbx), an epoetin alfa biosimilar approved for use in the United States, for intravenous or subcutaneous use.

The information provided in this document is intended for informational purposes only and is not a comprehensive description of potential coding requirements for RETACRIT. Coding and coverage policies change periodically and often without warning. The healthcare provider is solely responsible for determining coverage and reimbursement parameters and appropriate coding for treatment of his/her patients. The information provided should not be considered a guarantee of coverage or reimbursement for RETACRIT.

Please see Important Safety Information and Indications on pages 14-16, and full Prescribing Information, including BOXED WARNING and Medication Guide, available at RetacritHCP.com.



Making your patients' support needs a priority. Together.

At Pfizer Oncology Together, patient support is at the core of everything we do. We've gathered resources and developed tools to help patients and their loved ones throughout RETACRIT treatment. From helping to identify financial assistance options to connecting patients to resources for emotional support, your patients' needs are our priority.*



Benefits Verification

We can help determine a patient's coverage and out-of-pocket costs.

Prior Authorization (PA) Assistance

We can coordinate with a patient's insurer to determine the PA requirements. After a PA request is submitted, we can follow up with the payer until a final outcome is determined.

Appeals Assistance

We can review the reasons for a denied claim and provide information on payer requirements. After an appeal is submitted, we can follow up with the payer until a final outcome is determined.

Billing and Coding Assistance for Injectable Products

For your patient claim submissions, we provide easy access to sample forms and template letters, along with billing and coding information for physician office and hospital outpatient settings of care.

Patient Financial Assistance

We can help patients understand their benefits and connect them with financial assistance resources.



FOR LIVE, PERSONALIZED SUPPORT

Call **1-877-744-5675** (Monday–Friday 8 AM–8 PM ET)

VISIT

PfizerOncologyTogether.com

*Some services are provided through third-party organizations that operate independently and are not controlled by Pfizer. Availability of services and eligibility requirements are determined solely by these organizations.

Please see **Important Safety Information and Indications** on pages 14-16, and **full Prescribing Information, including BOXED WARNING and Medication Guide**, available at RetacritHCP.com.

INJECTION
Retacrit™
epoetin alfa-epbx
Pfizer Oncology

Pfizer
Oncology
together™

Coding Overview

In the physician office, hospital outpatient department, and dialysis sites of care, Medicare Administrative Contractors (MACs), private commercial payers, and Medicaid may recognize the following codes for reporting RETACRIT on claim forms. RETACRIT will be covered under the End-Stage Renal Disease (ESRD) Prospective Payment System for utilization in the dialysis setting.

Coding for RETACRIT

The Centers for Medicare & Medicaid Services (CMS) assigned RETACRIT product-specific Healthcare Common Procedure Coding System (HCPCS) codes to identify ESRD and non-ESRD utilization of RETACRIT. The HCPCS code used to report RETACRIT is different for ESRD and non-ESRD use. HCPs may use the following HCPCS codes for all payers in all settings of care.¹

HCPCS Code ¹	Descriptor
Q5105	Injection, epoetin alfa, biosimilar, (Retacrit) (for ESRD on dialysis), 100 units
Q5106	Injection, epoetin alfa, biosimilar, (Retacrit) (for non-ESRD use), 1,000 units

Modifiers may be included on ESRD and non-ESRD claims to provide additional information. The JW modifier is used to report the amount of the drug that is unused after administration to a patient. For Medicare and some payers, the unused amount should be reported on a separate line of the claim form, and the claim should include the drug code, modifier, and number of units discarded.² When reporting the administration of erythropoiesis-stimulating agents (ESAs) on non-ESRD claims, Medicare and some payers may require modifier EA, EB, or EC to specify anemia.³ Modifier ED or EE and GS may be used to describe hematocrit levels.⁴ For ESRD claims, some payers may require modifiers JA or JB to be reported, indicating the route of administration.⁵ Additional modifiers may also be considered appropriate when submitting claims.

HCPCS Modifier ¹	Descriptor
JW	Drug amount discarded/not administered to any patient
EA	ESA administered to treat anemia due to chemotherapy
EB	ESA administered to treat anemia due to radiotherapy
EC	ESA administered to treat anemia not due to radiotherapy or chemotherapy
ED	Hematocrit level has exceeded 39% (or hemoglobin level has exceeded 13.0 G/dL) for 3 or more consecutive billing cycles immediately prior to and including the current cycle
EE	Hematocrit level has not exceeded 39% (or hemoglobin level has not exceeded 13.0 G/dL) for 3 or more consecutive billing cycles immediately prior to and including the current cycle
EJ	Subsequent claims for a defined course of therapy, e.g., EPO, sodium hyaluronate, infliximab
GS	Dosage of erythropoietin stimulating agent has been reduced and maintained in response to hematocrit or hemoglobin level
JA	Administered intravenously
JB	Administered subcutaneously

Please see **Important Safety Information and Indications** on pages 14-16, and **full Prescribing Information, including BOXED WARNING and Medication Guide**, available at RetacritHCP.com.



RETACRIT National Drug Codes for Pfizer

National Drug Codes (NDCs) are unique 10-digit, 3-segment numbers used to identify drugs.⁶

Strength ⁷	Vial Size	10-Digit NDC
2,000 Units/mL	1 mL single-dose vial	0069-1305-01
3,000 Units/mL	1 mL single-dose vial	0069-1306-01
4,000 Units/mL	1 mL single-dose vial	0069-1307-01
10,000 Units/mL	1 mL single-dose vial	0069-1308-01
40,000 Units/mL	1 mL single-dose vial	0069-1309-01

NDC Conversion Example

For reimbursement purposes, some payers may require the HCP to include NDCs on the claim form. For claims-reporting purposes, some payers may also require HCPs to convert the 10-digit NDC to an 11-digit NDC by adding a “0” (zero) where appropriate to create a 5-4-2 configuration. The zero is added in front of the first segment of numbers when the 10-digit format is the 4-4-2 configuration. See placement of the red zero in the example below.

Strength	Vial Size	10-Digit NDC	11-Digit NDC
2,000 Units/mL	1 mL single-dose vial	0069-1305-01	<u>0</u> 0069-1305-01

Please see [Important Safety Information](#) and [Indications](#) on pages 14-16, and [full Prescribing Information, including BOXED WARNING and Medication Guide](#), available at RetacritHCP.com.

Coding for RETACRIT Administration Services

Current Procedural Terminology (CPT®) codes define specific medical procedures performed by physicians.⁸

The following codes may be used to report the administration of RETACRIT:

Type of Code	Code/Descriptor	Relevant Sites of Service
Administration: CPT® codes ⁸	96372: Therapeutic, prophylactic, or diagnostic injection (specify substance or drug); subcutaneous or intramuscular	
	96374: Therapeutic, prophylactic, or diagnostic injection (specify substance or drug); IV push, single or initial substance/drug (Used for a short IV infusion lasting 15 min or less)	Physician office, hospital outpatient department, dialysis facility
	96375: Therapeutic, prophylactic, or diagnostic injection (specify substance or drug); each additional sequential IV push of a new substance/drug (list separately in addition to code for primary procedure) (Used for a short IV infusion lasting 15 min or less)	

Hospital outpatient departments and dialysis facilities use revenue codes to report specific accommodations and/or ancillary charges.⁹

Type of Code	Code/Descriptor	Relevant Sites of Service
Revenue codes ¹⁰	0634: Drugs requiring specific identification – EPO under 10,000 units (For RETACRIT)	Hospital outpatient department, dialysis facility
	0635: Drugs requiring specific identification – EPO 10,000 units or more (For RETACRIT)	
	0636: Drugs requiring specific identification – detailed coding (For RETACRIT)	Hospital outpatient department
	0510: Clinic – general classification (For IV injection administered in the clinic)	

Key: CPT – Current Procedural Terminology; IV – intravenous; EPO – erythropoietin.
Current Procedural Terminology (CPT®) is a registered trademark of the American Medical Association.

Please see **Important Safety Information and Indications** on pages 14-16, and **full Prescribing Information, including BOXED WARNING and Medication Guide**, available at RetacritHCP.com.

Diagnosis Coding for RETACRIT

RETACRIT (epoetin alfa-epbx) is an FDA-approved biosimilar.

The International Classification of Diseases, 10th Revision, Clinical Modification (ICD-10-CM) code set should be used, as appropriate, to report the patient-specific diagnosis.

RETACRIT is typically reported using a primary diagnosis code for anemia and a secondary diagnosis code for a disease indication. Payer-specific coding requirements should be verified by the HCP, including the order (eg, primary, secondary, etc) of required codes. HCPs should verify payer-specific coding requirements before submitting a claim, as these may vary by payer.

Coding to Report Anemia and Related Conditions

ICD-10-CM codes to report anemia and related conditions may include, but are not limited to, the following codes:

ICD-10-CM Code ¹¹	Code Descriptor
B20	Human immunodeficiency virus disease (Code first human immunodeficiency virus [HIV] disease complicating pregnancy, childbirth, and the puerperium, if applicable [O98.7-]. Use additional code[s] to identify all manifestations of HIV infection)
D63.1*	Anemia in chronic kidney disease (Code first underlying chronic kidney disease [N18.-])
D64.81	Anemia due to antineoplastic chemotherapy
I12.0	Hypertensive chronic kidney disease with stage 5 chronic kidney disease or end-stage renal disease (Use additional code to identify the stage of chronic kidney disease [N18.5, N18.6])
N18.3	Chronic kidney disease, stage 3 (moderate)
N18.4	Chronic kidney disease, stage 4 (severe)
N18.5	Chronic kidney disease, stage 5
Z21	Asymptomatic human immunodeficiency virus infection status (Code first human immunodeficiency virus disease complicating pregnancy, childbirth, and the puerperium, if applicable [O98.7-])

Key: ICD-10-CM – International Classification of Diseases, 10th Revision, Clinical Modification.

*Only to be billed by a nephrologist.¹²

Please see **Important Safety Information and Indications** on pages 14-16, and **full Prescribing Information, including BOXED WARNING and Medication Guide**, available at RetacritHCP.com.

RETACRIT Billing Units

ESRD

The RETACRIT HCPCS code Q5105 is described as “Injection, epoetin alfa, biosimilar, (Retacrit) (for ESRD on dialysis), 100 units.” Each dose increment of 100 Units equals 1 billing unit. For example, a 2,000 Units/mL vial of RETACRIT represents 20 billing units of Q5105. See the chart below correlating a vial of RETACRIT administered with the number of billing units based on the description of Q5105.

Strength	Vial Size	Number of Q5105 Billing Units (100 Units epoetin alfa-epbx biosimilar per 1 billing unit) Equivalent to the Units of RETACRIT in Each Vial
2,000 Units/mL	1 mL single-dose vial	20 units
3,000 Units/mL	1 mL single-dose vial	30 units
4,000 Units/mL	1 mL single-dose vial	40 units
10,000 Units/mL	1 mL single-dose vial	100 units
40,000 Units/mL	1 mL single-dose vial	400 units

Non-ESRD

The RETACRIT HCPCS code Q5106 is described as “Injection, epoetin alfa, biosimilar, (Retacrit) (for non-ESRD on dialysis) 1,000 Units.” Each dose increment of 1,000 Units equals 1 billing unit. For example, a 2,000 Units/mL vial of RETACRIT represents 2 billing units of Q5106. See the chart below correlating a vial of RETACRIT administered with the number of billing units based on the description of Q5106.

Strength	Vial Size	Number of Q5106 Billing Units (1,000 Units epoetin alfa-epbx biosimilar per 1 billing unit) Equivalent to the Units of RETACRIT in Each Vial
2,000 Units/mL	1 mL single-dose vial	2 units
3,000 Units/mL	1 mL single-dose vial	3 units
4,000 Units/mL	1 mL single-dose vial	4 units
10,000 Units/mL	1 mL single-dose vial	10 units
40,000 Units/mL	1 mL single-dose vial	40 units

Please see [Important Safety Information](#) and [Indications](#) on pages 14-16, and [full Prescribing Information, including BOXED WARNING and Medication Guide](#), available at RetacritHCP.com.

Claims Submission Checklist

The following may be considered to assist with submitting claims completely and accurately, which is important for timely claims processing, for appropriate payment, and to avoid denied claims.



- ✓ Provide the patient name, address, and insurance identification number, and review these for accuracy
- ✓ Include the HCP's name, National Provider Identifier (NPI), and payer-specific provider ID (if applicable)
- ✓ Indicate the appropriate place of service code (2-digit code) for where the treatment was provided
- ✓ Check to ensure that ICD-10-CM diagnosis codes, CPT procedure codes, and modifiers (if applicable) are consistent with information included in the patient's medical record
- ✓ Review the RETACRIT-specific information (eg, name of drug, HCPCS code, NDC, number of units, route and frequency of administration)

Please see Important Safety Information and Indications on pages 14-16, and full Prescribing Information, including BOXED WARNING and Medication Guide, available at RetacritHCP.com.

Sample Claim Form: CMS-1500, Physician Office Setting (Non-ESRD)

HEALTH INSURANCE CLAIM FORM
APPROVED BY NATIONAL UNIFORM CLAIM COMMITTEE (NUCC) 02/12

Item 19: If additional information is required to describe RETACRIT (eg, NDC), this information may be captured in Item 19

Item 21: Specify appropriate ICD-10-CM diagnosis code(s)

Item 21: Identify the type of ICD diagnosis code used (eg, enter a "0" for ICD-10-CM)

Item 24D: Specify appropriate HCPCS and CPT codes and modifiers; for example:

- Drug: Q5106 for RETACRIT
- Administration: 96xxx for administration

Item 24E: Enter reference to the diagnosis for the CPT and HCPCS codes from Item 21

This sample form is intended as a reference for the coding and billing of RETACRIT. This form is not intended to be directive and the use of the recommended codes does not guarantee reimbursement. HCPs may deem other codes or policies more appropriate and should select the coding options that most accurately reflect their internal guidelines, payer requirements, practice patients, and services rendered.

Please see **Important Safety Information and Indications** on pages 14-16, and **full Prescribing Information, including BOXED WARNING and Medication Guide**, available at RetacritHCP.com.

Sample Claim Form: UB-04, Hospital Outpatient Setting (Non-ESRD)

This sample form is intended as a reference for the coding and billing of RETACRIT. This form is not intended to be directive and the use of the recommended codes does not guarantee reimbursement. HCPs may deem other codes or policies more appropriate and should select the coding options that most accurately reflect their internal guidelines, payer requirements, practice patients, and services rendered.

Form Locator (FL) 44: Specify appropriate HCPCS and CPT codes and modifiers; for example:

- Drug: Q5106 for RETACRIT
- Administration: 96xxx for drug administration

FL 39-41: Enter value code 48 with **hemoglobin** level reading or value code 49 with **hematocrit** level reading

FL 46: Specify the billing units. For example, 1 billing unit = **1,000** Units of epoetin alfa-epbx biosimilar (RETACRIT) for HCPCS code Q5106. To bill 2,000 Units of RETACRIT, enter 2 billing units. To bill 1 96xxx for drug administration, enter 1 billing unit

FL 42 and 43: Specify revenue codes and describe procedure; for example:

- 0636: Drugs requiring specific identification – detailed coding (For RETACRIT)
- 0510: Clinic – general classification (For IV injection administered in hospital-based dialysis clinics)

Note: Other revenue codes may apply

FL 67: Specify appropriate ICD-10-CM diagnosis code(s)

FL 66: Identify the type of ICD diagnosis code used (eg, enter a “0” for ICD-10-CM)

Please see **Important Safety Information and Indications** on pages 14-16, and **full Prescribing Information, including BOXED WARNING and Medication Guide**, available at RetacritHCP.com.

Sample Claim Form: UB-04, Dialysis Provider for Use in Dialysis (ESRD)

This sample form is intended as a reference for the coding and billing of RETACRIT. This form is not intended to be directive and the use of the recommended codes does not guarantee reimbursement. HCPs may deem other codes or policies more appropriate and should select the coding options that most accurately reflect their internal guidelines, payer requirements, practice patients, and services rendered.

FL 44: Specify appropriate HCPCS and CPT codes and modifiers; for example:

- Drug: Q5105 for RETACRIT
- Administration: 96xxx for drug administration

FL 39-41: Enter value code 48 with **hemoglobin** level reading or value code 49 with **hematocrit** level reading

FL 46: Specify the billing units. For example, 1 billing unit = **100** Units of epoetin alfa-epbx biosimilar (RETACRIT) for HCPCS code Q5105. To bill 2,000 Units of RETACRIT, enter 20 billing units. To bill 1 96xxx for drug administration, enter 1 billing unit

FL 42 and 43: Specify revenue codes and describe procedures, for example:

- 0634: Drugs requiring specific identification – EPO under 10,000 units (For RETACRIT), or
- 0635: Drugs requiring specific identification – EPO 10,000 units or more (For RETACRIT)
- 0510: Clinic – general classification (For IV injection administered in hospital-based dialysis clinics)

Note: Other revenue codes may apply

FL 67: Specify appropriate ICD-10-CM diagnosis code(s)

FL 66: Identify the type of ICD diagnosis code used (eg, enter a “0” for ICD-10-CM)

Please see **Important Safety Information and Indications** on pages 14-16, and **full Prescribing Information, including BOXED WARNING and Medication Guide**, available at RetacritHCP.com.

References

1. Centers for Medicare & Medicaid Services (CMS). 2019 Alpha-numeric HCPCS file. <https://www.cms.gov/Medicare/Coding/HCPCSReleaseCodeSets/Downloads/2019-Alpha-Numeric-HCPCS-File.zip>. Accessed May 17, 2019.
2. Centers for Medicare & Medicaid Services (CMS). Medicare program JW modifier: drug/biological amount discarded/not administered to any patient frequently asked questions. <https://www.cms.gov/Medicare/Medicare-Fee-for-Service-Payment/HospitalOutpatientPPS/Downloads/JW-Modifier-FAQs.pdf>. Accessed May 20, 2019.
3. Centers for Medicare & Medicaid Services (CMS). Pub 100-04 Medicare claims processing: reporting of hematocrit or hemoglobin levels on all claims for the administration of erythropoiesis stimulating agents (ESAs), implementation of new modifiers for non-ESRD indications, and reporting of hematocrit/hemoglobin levels on all non-ESRD, non-ESA claims requesting payment for anti-anemia drugs. <https://www.cms.gov/Regulations-and-Guidance/Guidance/Transmittals/downloads/R1412CP.pdf>. Accessed May 20, 2019.
4. Centers for Medicare & Medicaid Services (CMS). MLN Matters Number: MM5700: modification to the national monitoring policy for erythropoiesis stimulating agents (ESAs) for end-stage renal disease (ESRD) patients treated in renal dialysis facilities. <https://www.cms.gov/Outreach-and-Education/Medicare-Learning-Network-MLN/MLNMattersArticles/downloads/MM5700.pdf>. Accessed May 20, 2019.
5. Centers for Medicare & Medicaid Services. Medicare Learning Network (MLN) Matters. Implementation of the Medicare Improvements for Patients and Providers Act of 2008 (MIPPA) 153c End Stage Renal Disease (ESRD) Quality Incentive Program (QIP) and other requirements for ESRD claims. <https://www.cms.gov/Outreach-and-Education/Medicare-Learning-Network-MLN/MLNMattersArticles/Downloads/MM7460.pdf>. Accessed May 20, 2019.
6. U.S. Food and Drug Administration (FDA). National Drug Code directory. <https://www.fda.gov/drugs/informationondrugs/ucm142438.htm>. Accessed May 20, 2019.
7. RETACRIT [package insert]. New York, NY: Pfizer Inc.; 2019.
8. American Medical Association. 2019 CPT Professional Edition. Current Procedural Terminology (CPT®) is copyright 2018 by the American Medical Association. All rights reserved. Chicago, IL: AMA; 2018.
9. Centers for Medicare & Medicaid Services. Transmittal 167. <https://www.cms.gov/Regulations-and-Guidance/Guidance/Transmittals/downloads/R167CP.pdf>. Accessed May 20, 2019.
10. Research Data Assistance Center (ResDAC). Revenue Center Code. <https://www.resdac.org/sites/resdac.umn.edu/files/Revenue%20Center%20Table.txt>. Accessed May 20, 2019.
11. 2019 International Statistical Classification of Diseases and Related Health Problems, 10th Revision, Clinical Modification (ICD-10-CM) Expert for Hospitals. Copyright 2018 Optum360, LLC. All rights reserved.
12. Centers for Medicare & Medicaid Services. CMS Manual. SUBJECT: ICD-10 and other coding revisions to national coverage determinations (NCDs). <https://www.cms.gov/Regulations-and-Guidance/Guidance/Transmittals/2017Downloads/R19750TN.pdf>. Accessed May 20, 2019.

Please see **Important Safety Information** and **Indications** on pages 14-16, and **full Prescribing Information, including BOXED WARNING and Medication Guide**, available at RetacritHCP.com.



IMPORTANT SAFETY INFORMATION

WARNINGS: ESAs INCREASE THE RISK OF DEATH, MYOCARDIAL INFARCTION, STROKE, VENOUS THROMBOEMBOLISM, THROMBOSIS OF VASCULAR ACCESS AND TUMOR PROGRESSION OR RECURRENCE

CHRONIC KIDNEY DISEASE:

- In controlled, trials patients experienced greater risks for death, serious adverse cardiovascular reactions, and stroke when administered erythropoiesis-stimulating agents (ESAs) to target a hemoglobin level of greater than 11 g/dL.
- No trial has identified a hemoglobin target level, ESA dose, or dosing strategy that does not increase these risks.
- Use the lowest RETACRIT™ dose sufficient to reduce the need for red blood cell (RBC) transfusions.

CANCER:

- ESAs shortened overall survival and/or increased the risk of tumor progression or recurrence in clinical studies of patients with breast, non-small cell lung, head and neck, lymphoid, and cervical cancers.
- To decrease these risks, as well as the risk of serious cardiovascular and thromboembolic reactions, use the lowest dose needed to avoid RBC transfusions.
- Use ESAs only for anemia from myelosuppressive chemotherapy.
- ESAs are not indicated for patients receiving myelosuppressive chemotherapy when the anticipated outcome is cure.
- Discontinue following the completion of a chemotherapy course.

PERISURGERY:

- Due to increased risk of deep venous thrombosis (DVT), DVT prophylaxis is recommended.

CONTRAINDICATIONS

RETACRIT™ is contraindicated in patients with:

- Uncontrolled hypertension.
- Pure red cell aplasia (PRCA) that begins after treatment with RETACRIT™ or other erythropoietin protein drugs.
- Serious allergic reactions to RETACRIT™ or other epoetin alfa products.

Please see **full Prescribing Information, including BOXED WARNING and Medication Guide, available at RetacritHCP.com.**

INCREASED MORTALITY, MYOCARDIAL INFARCTION, STROKE, AND THROMBOEMBOLISM

- In controlled clinical trials of patients with CKD comparing higher hemoglobin targets (13 - 14 g/dL) to lower targets (9 - 11.3 g/dL), epoetin alfa increased the risk of death, myocardial infarction, stroke, congestive heart failure, thrombosis of hemodialysis vascular access, and other thromboembolic events in the higher target groups.
- Using ESAs to target a hemoglobin level of greater than 11 g/dL increases the risk of serious adverse cardiovascular reactions and has not been shown to provide additional benefit. Use caution in patients with coexistent cardiovascular disease and stroke. Patients with CKD and an insufficient hemoglobin response to ESA therapy may be at even greater risk for cardiovascular reactions and mortality than other patients. A rate of hemoglobin rise of greater than 1 g/dL over 2 weeks may contribute to these risks.
- In controlled clinical trials of patients with cancer, epoetin alfa increased the risks for death and serious adverse cardiovascular reactions. These adverse reactions included myocardial infarction and stroke.
- In controlled clinical trials, ESAs increased the risk of death in patients undergoing coronary artery bypass graft surgery (CABG) and the risk of deep venous thrombosis (DVT) in patients undergoing orthopedic procedures.

INCREASED MORTALITY AND/OR INCREASED RISK OF TUMOR PROGRESSION OR RECURRENCE IN PATIENTS WITH CANCER

- ESAs resulted in decreased locoregional control/progression-free survival (PFS) and/or overall survival (OS). Adverse effects on PFS and/or OS were observed in studies of patients receiving chemotherapy for breast cancer, lymphoid malignancy, and cervical cancer; in patients with advanced head and neck cancer receiving radiation therapy; and in patients with non-small cell lung cancer or various malignancies who were not receiving chemotherapy or radiotherapy.

HYPERTENSION

- RETACRIT™ is contraindicated in patients with uncontrolled hypertension. Following initiation and titration of epoetin alfa, approximately 25 % of patients on dialysis required initiation of or increases in antihypertensive therapy; hypertensive encephalopathy and seizures have been reported in patients with CKD receiving RETACRIT™.

Continued on the next page

IMPORTANT SAFETY INFORMATION (Continued)

- Appropriately control hypertension prior to initiation of and during treatment with RETACRIT™. Reduce or withhold RETACRIT™ if blood pressure becomes difficult to control. Advise patients of the importance of compliance with antihypertensive therapy and dietary restrictions.

SEIZURES

- Epoetin alfa products, including RETACRIT™, increase the risk of seizures in patients with CKD. During the first several months following initiation of RETACRIT™, monitor patients closely for premonitory neurologic symptoms. Advise patients to contact their healthcare practitioner for new-onset seizures, premonitory symptoms or change in seizure frequency.

LACK OR LOSS OF HEMOGLOBIN RESPONSE TO RETACRIT™

- For lack or loss of hemoglobin response to RETACRIT™, initiate a search for causative factors (e.g., iron deficiency, infection, inflammation, bleeding). If typical causes of lack or loss of hemoglobin response are excluded, evaluate for PRCA. In the absence of PRCA, follow dosing recommendations for management of patients with an insufficient hemoglobin response to RETACRIT™ therapy.

PURE RED CELL APLASIA

- Cases of PRCA and of severe anemia, with or without other cytopenias that arise following the development of neutralizing antibodies to erythropoietin have been reported in patients treated with epoetin alfa. This has been reported predominantly in patients with CKD receiving ESAs by subcutaneous administration. PRCA has also been reported in patients receiving ESAs for anemia related to hepatitis C treatment (an indication for which RETACRIT™ is not approved).
- If severe anemia and low reticulocyte count develop during treatment with RETACRIT™, withhold RETACRIT™ and evaluate patients for neutralizing antibodies to erythropoietin. **Contact Pfizer Inc. at 1-800-438-1985 to perform assays for binding and neutralizing antibodies.** Permanently discontinue RETACRIT™ in patients who develop PRCA following treatment with RETACRIT™ or other erythropoietin protein drugs. Do not switch patients to other ESAs.

SERIOUS ALLERGIC REACTIONS

- Serious allergic reactions, including anaphylactic reactions, angioedema, bronchospasm, skin rash, and urticaria may occur with epoetin alfa products. Immediately and permanently discontinue RETACRIT™ and administer appropriate therapy if a serious allergic or anaphylactic reaction occurs.

SEVERE CUTANEOUS REACTIONS

- Blistering and skin exfoliation reactions, including Erythema multiforme and Stevens-Johnson Syndrome (SJS)/Toxic Epidermal Necrolysis (TEN), have been reported in patients treated with ESAs (including epoetin alfa) in the postmarketing setting. Discontinue RETACRIT™ therapy immediately if a severe cutaneous reaction, such as SJS/TEN, is suspected.

RISK IN PATIENTS WITH PHENYLKETONURIA

- Phenylalanine can be harmful to patients with phenylketonuria (PKU). RETACRIT™ contains phenylalanine, a component of aspartame. Each 1 mL single-dose vial of 2,000, 3,000, 4,000, 10,000, and 40,000 Units of epoetin alfa-epbx injection contains 0.5 mg of phenylalanine. Before prescribing RETACRIT™ to a patient with PKU, consider the combined daily amount of phenylalanine from all sources, including RETACRIT™.

DIALYSIS MANAGEMENT

- Patients may require adjustments in their dialysis prescriptions after initiation of RETACRIT™. Patients receiving RETACRIT™ may require increased anticoagulation with heparin to prevent clotting of the extracorporeal circuit during hemodialysis.

ANEMIA IN PATIENTS WITH CHRONIC KIDNEY DISEASE

- Adverse reactions in ≥5 % of epoetin alfa-treated patients on dialysis were hypertension, arthralgia, muscle spasm, pyrexia, dizziness, medical device malfunction, vascular occlusion and upper respiratory tract infection.

ANEMIA DUE TO CHEMOTHERAPY IN PATIENTS WITH CANCER

- Adverse reactions in ≥5 % of epoetin alfa-treated patients in clinical studies were nausea, vomiting, myalgia, arthralgia, stomatitis, cough, weight decrease, leukopenia, bone pain, rash, hyperglycemia, insomnia, headache, depression, dysphagia, hypokalemia, and thrombosis.

Please see **full Prescribing Information, including BOXED WARNING and Medication Guide, available at RetacritHCP.com.**

Continued on the next page

IMPORTANT SAFETY INFORMATION (Continued)

SURGERY/PERISURGERY

- Adverse reactions in $\geq 5\%$ of epoetin alfa-treated patients in clinical studies were nausea, vomiting, pruritus, headache, injection site pain, chills, deep vein thrombosis, cough, and hypertension.

ANEMIA DUE TO ZIDOVUDINE IN PATIENTS WITH HIV-INFECTION

- Adverse reactions in $\geq 5\%$ of epoetin alfa-treated patients in clinical studies were pyrexia, cough, rash, and injection site irritation.

INDICATIONS AND LIMITATIONS OF USE

ANEMIA DUE TO CHRONIC KIDNEY DISEASE

RETACRIT™ is indicated for the treatment of anemia due to chronic kidney disease (CKD), including patients on dialysis and not on dialysis, to decrease the need for red blood cell (RBC) transfusion.

ANEMIA DUE TO ZIDOVUDINE IN PATIENTS WITH HIV-INFECTION

RETACRIT™ is indicated for the treatment of anemia due to zidovudine administered at $\leq 4,200$ mg/week in patients with HIV-infection with endogenous serum erythropoietin levels of ≤ 500 mUnits/mL.

ANEMIA DUE TO CHEMOTHERAPY IN PATIENTS WITH CANCER

RETACRIT™ is indicated for the treatment of anemia in patients with nonmyeloid malignancies where anemia is due to the effect of concomitant myelosuppressive chemotherapy, and upon initiation, there is a minimum of two additional months of planned chemotherapy.

REDUCTION OF ALLOGENEIC RED BLOOD CELL TRANSFUSIONS IN PATIENTS UNDERGOING ELECTIVE, NONCARDIAC, NONVASCULAR SURGERY

RETACRIT™ is indicated to reduce the need for allogeneic RBC transfusions among patients with perioperative hemoglobin > 10 to ≤ 13 g/dL who are at high risk for perioperative blood loss from elective, noncardiac, nonvascular surgery. RETACRIT™ is not indicated for patients who are willing to donate autologous blood preoperatively.

RETACRIT™ has not been shown to improve quality of life, fatigue, or patient well-being.

RETACRIT™ is not indicated for use:

- In patients with cancer receiving hormonal agents, biologic products, or radiotherapy, unless also receiving concomitant myelosuppressive chemotherapy.
- In patients with cancer receiving myelosuppressive chemotherapy when the anticipated outcome is cure.
- In patients with cancer receiving myelosuppressive chemotherapy in whom the anemia can be managed by transfusion.
- In patients scheduled for surgery who are willing to donate autologous blood.
- In patients undergoing cardiac or vascular surgery.
- As a substitute for RBC transfusions in patients who require immediate correction of anemia.

Please see **full Prescribing Information, including BOXED WARNING and Medication Guide, available at RetacritHCP.com.**

PP-RET-USA-0183 © 2019 Pfizer Inc. All rights reserved. August 2019

